

ON GROTHENDIECK SUBSPACES

S. S. KUTATELADZE

February 21, 2005

Let X be a Riesz space, and let Y be a Kantorovich (or Dedekind complete Riesz) space Y with base a complete Boolean algebra B . In [1] we have described every order bounded operator T from X to Y that may be presented as the difference of some Riesz homomorphisms. An operator T is such a difference if and only if the kernel of the stratum bT of T is a Riesz subspace of X for all $b \in B$. We will give an analogous description for an order bounded operator T whose modulus may be presented as the sum of two Riesz homomorphisms in terms of the properties of the strata of T .

Note that the sums of Riesz homomorphisms were first described by S. J. Bernau, C. B. Huijsmans, and B. de Pagter in terms of n -disjoint operators in [2]. A survey of some conceptually close results on n -disjoint operators is given in [3, §5.6]. In this note we reveal the connection between the 2-disjoint operators and Grothendieck subspaces.

Theorem. *The modulus of an order bounded operator $T : X \rightarrow Y$ is the sum of some pair of Riesz homomorphisms if and only if the kernel of each stratum bT of T with $b \in B$ is a Grothendieck subspace of the ambient Riesz space X .*

Recall that a subspace H of a Riesz space is a G -space or *Grothendieck subspace* provided that H enjoys the following property:

$$(1) \quad (\forall x, y \in H) \ (x \vee y \vee 0 + x \wedge y \wedge 0 \in H).$$

The history of (1) is as follows: In 1955 A. Grothendieck distinguished the subspaces that satisfy (1) in the space $C(Q, \mathbb{R})$ of continuous real functions on a compact space Q (cp. [4]). He determined such a subspace as the set of functions f satisfying some family A of relations of the form

$$f(q_\alpha^1) = \lambda_\alpha f(q_\alpha^2) \ (q_\alpha^1, q_\alpha^2 \in Q; \ \lambda_\alpha \in \mathbb{R}, \ \alpha \in A).$$

These subspaces due to A. Grothendieck gave examples of the L_1 -predual Banach spaces other than AM -spaces. In 1969 J. Lindenstrauss and D. Wulpert characterized the subspaces by using (1) and introduced the concept of G -space (cp. [5]). Some related properties of Grothendieck spaces are presented also in [6] and [7].

As in [1], we rest the proof of the above-stated theorem on the technique of “non-standard scalarization” reducing operator problems to the case of functionals. Using the facts of Riesz space theory and Boolean valued analysis we will presume the terminology and notation of [8].

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

We argue further as follows: Using the functors of canonical embedding and descent to the Boolean valued universe $\mathbb{V}^{(B)}$, we reduce the matter to characterizing a Grothendieck hyperspace that serves as the kernel of an order bounded functional over a dense subring of the reals $\mathbb{R}$. The scalar case is settled by the following four lemmas.

Lemma 1. *A functional l is the sum of some pair of Riesz homomorphisms if and only if l is a positive functional with kernel a Grothendieck subspace.*

Lemma 2. *Given an order bounded functional l on a vector lattice X , assume that $l_+ \neq 0$ and $l_- \neq 0$. The kernel $\ker(l)$ is a Grothendieck subspace of if and only if l_+ and l_- are Riesz homomorphisms on X (or, which is the same, $\ker(l)$ is a Riesz subspace).*

Lemma 3. *The kernel of an order bounded functional is a Grothendieck subspace if and only if so is the kernel of the modulus of this functional.*

Lemma 4. *The kernel of an order bounded functional is a Grothendieck subspace if and only if the modulus of this functional is the sum of a pair of Riesz homomorphisms.*

REFERENCES

1. Kutateladze S. S., *On differences of Riesz homomorphisms*, Sibirsk. Mat. Zh. **46** (2005), no. 2.
2. Bernau S. J., Huijsmans C.B., and de Pagter B., *Sums of lattice homomorphisms*, Proc. Amer. Math. Soc. **115** (1992), no. 1, 151–156.
3. Kusraev A. G., *Dominated Operators*, Kluwer Academic Publishers, Dordrecht, 2000.
4. Grothendieck A., *Une caractérisation vectorielle-métrique des espaces L^1* , Canad. J. Math. **4** (1955), 552–561.
5. Lindenstrauss J. and Wulbert D. E., *On the classification of the Banach spaces whose duals are L_1 -spaces*, J. Funct. Anal. **4** (1969), no. 3, 332–349.
6. Semadeni Zb., *Banach Spaces of Continuous Functions*, Polish Scientific Publishers, Warszawa, 1971.
7. Lacey H. E., *The Isometric Theory of Classical Banach Spaces*, Springer-Verlag, Berlin etc., 1973.
8. Kusraev A. G. and Kutateladze S. S., *Boolean Valued Analysis*, Kluwer Academic Publishers, Dordrecht, 1999.

NOVOSIBIRSK